



SPECIMEN LABEL

MCPA AMIINE 4

**HERBICIDE FOR SELECTIVE CONTROL OF CERTAIN WEEDS
IN WHEAT, OATS, BARLEY, RYE, RICE, FLAX,
ESTABLISHED GRASSLANDS AND NON-CROP AREAS**

ACTIVE INGREDIENT:	
Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid*	48.89%
INERT INGREDIENTS:	51.11%
TOTAL	100.00%

*This product contains 3.7 pounds 2-methyl-4-chlorophenoxyacetic acid per gallon or 39.9% Isomer specific by AOAC Method No. 6.A-18-22 (13th Ed.)

EPA Reg. No. 42750-14

EPA Est. No. 42750-MO-1

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

See additional Precautionary Statements on left side panel of page two.

Manufactured by:
ALBAUGH, INC.
121 N.E. 18th Street
Ankeny, Iowa 50021

NET CONTENTS
____ GALS.
____ LITERS

**FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE
CALL CHEMTREC (800) 424-9300**

Before buying or using this product, read “Warranty Limitations and Disclaimer” elsewhere on this label. If terms are not acceptable, return **unopened** package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Warranty Limitations and Disclaimer.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed, absorbed through skin or inhaled. Do not get in eyes, or on clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product’s concentrate. Do not reuse them. Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Drift or run-off may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Avoid use of small-diameter spray nozzles. Do not apply when weather conditions favor drift from target area, as this product may injure cotton, beans, other vegetables, certain legumes and ornamentals.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. Do not store below 40° F or expose to subfreezing temperatures. Handle in accordance with information given under “Precautionary Statements.” In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust or other absorbent material. Scrape up and dispose of in accordance with information given under “Disposal.” Repackage and relabel useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

DISPOSAL

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

METAL CONTAINERS - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

PLASTIC CONTAINERS - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS

Under no circumstances should MCPA Amine Herbicide be used in the vicinity of vegetables, flowers, grapes, tomatoes, cotton, beans and other legumes, etc., or other susceptible crops as severe damage may result. Extreme care must be exercised to prevent drifting of this material. Do not apply on windy days. Coarse sprays are less likely to drift. Do not apply this product through any type of irrigation system. Do not use around the home, recreation areas or similar sites. Equipment used to apply MCPA Amine should not be used to apply other agricultural chemicals to susceptible crops. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

AMOUNT OF SPRAY TO APPLY

Apply 5 to 15 gallons of total spray per acre when making applications with ground equipment and 3 to 5 gallons of total spray per acre when making applications by aircraft, unless directed otherwise under specific directions. Carefully read the Environmental Hazards section of the Precautionary Statements for further information on spray volume.

CONTROLS THESE WEEDS SUSCEPTIBLE (MOST AREAS)

Cocklebur	Marshelder	Stinkweed
Dragonhead Mint	Mustards	Wild Radish
Fanweed	(Annual)	Yellow Rocket
Field Peppergrass	Pennycress	and many others
Goatsbeard	or Wintercress	
Lambsquarters	Puncturevine	
(Goosefoot)	Shepherdspurse	

LESS SUSCEPTIBLE

Buttercups	Plantains	Vetch
Canada Thistle	Purslane	White Top
Dandelions	Russian Pigweed	and Hoary Cress
Hempnettle	Sow Thistle	and many others
Kochia	Stinging Nettle	
Pigweed	Sunflower	

DIRECTIONS FOR USE

SELECTIVE SPRAYING: Wheat, Oats, Barley and Rye - Apply as a water mix spray by ground sprayer or airplane. Use 1/2 to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Use up to 3 pints per acre for less susceptible weeds after crop has tillered and up to early boot stage. Do not spray from boot to dough stage. For small grains underseeded with legumes, see the following paragraph.

SMALL GRAINS: Underseeded with Alfalfa, Lespedeza, Red and White Clover: For emergency control of serious infestations of Mustard, Yellow Rocket and other susceptible broadleaf weeds, apply 1/4 to 1/2 pint in not more than 5 to 10 gallons of water per acre. Make application after cereal is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy which, together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. Do not apply to small grains underseeded with Vetch or Sweet Clover, which are very susceptible. There is also some risk to other legumes if only thinly protected by a canopy. Do not forage or graze meat animals on treated areas within 7 days of slaughter.

RED CLOVER AND ALFALFA (for use in the Pacific Northwest only): For control of Yellow Rocket and other susceptible annual weeds such as Pennycress and Fanweed, use 1 pint per acre of MCPA AMINE. Only use 1/2 pint per acre on new stands after Clover has two or more true leaves. Old stands of Red Clover may be retarded by application of MCPA. Apply late in fall following frosts when the legumes are dormant before active growth starts. The temperature at the time of spraying should be above 40 F.

CANNING PEAS (for use in the Pacific Northwest only): Use MCPA Amine at 1/4 to 3/4 pint in sufficient water to cover 1 acre. Apply to peas after the 3 node stage and before the first pea flowering. Do not apply during bloom period of crop. To control Canada Thistle, use 1/2 to 3/4 pint of MCPA Amine per acre. Peas may be injured somewhat at the higher rate of application, but if thistle growth is heavy, control will more than compensate for injury to peas.

FLAX: Use 1/4 to 1/2 pint. Apply by ground sprayer or by airplane. Use sufficient water to give uniform and adequate coverage. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada Thistle is present, it may be necessary to go as high as 3/4 pint per acre to prevent seed head production. Some injury to the flax may result. Do not forage or graze meat animals on treated areas within 7 days of slaughter.

ESTABLISHED GRASSLAND AND PASTURES: Use 1 to 3 pints per acre in sufficient water (10 to 100 gallons) in airplane or ground sprayer application and give thorough coverage. Use higher rate for White Top, Canada Thistle and other less easy-to-kill weeds; spray perennials in early bud to full bloom stage and also regrowth in fall. Spray other weeds in spring or fall. Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.

GRASSES GROWN FOR SEED: Use 1 to 2 pints per acre in sufficient water to give adequate coverage. Use higher rate where weed stands are heavy. In established grasses, apply in spring before head comes into boot and on seedling grass after grass has tillered.

NOTE: In weed control in grasses, repeat treatment may be needed for less susceptible weeds. White Clover and other legumes may be temporarily injured or killed. In some areas, Bent, Buffalo, Carpet and St. Augustine grasses may also be injured by the treatment.

RANGE AND TIMBERLANDS: For control of Whitebrush - Use 1-1/4 quarts in a mixture of 1 gallon of diesel oil and sufficient water to make 8 gallons of solution per acre. Apply in spring or fall under good moisture conditions, full leaf, before blossoms begin to fall.

NON-CROP SPRAYING

CANADA THISTLE, WHITE TOP AND MEADOW BUTTERCUP (In non-crop areas such as Roadsides, Fence Rows, Rights-of-Way and similar places): For spot treatment, use 1/4 pint to 3 to 4 gallons of water, or 3/4 gallon per acre in sufficient water to give coverage for most extensive areas. Spray

to wet weeds thoroughly when in bud to early bloom and again on fall regrowth. Do not forage or graze livestock or dairy animals on treated areas within 7 days of treatment. Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

WARRANTY LIMITATIONS AND DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller. When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing, of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of the seller will be one of the following, at the election of the seller:

1. Refund of purchase price paid by buyer or user for product bought or
2. Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of the Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.

Manufactured by:

ALBAUGH, INC
121 N.E. 18th Street, Ankeny, Iowa 50021

Rev. 10/99